

t3_mesfunc6
(TMF6RxLEucbjzhNdv1NiSNdyubb1Btk7ZDT)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k11_mesfunc1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v2_valued_0 : \iota \Rightarrow o$ be given. Let $k12_supinf_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v2_membered : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v3_valued_0 X0))) \Rightarrow (k1_seq_1 X0 X1 = k1_funct_1 X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v2_valued_0 X0))) \Rightarrow (k12_supinf_2 X0 X1 = k1_funct_1 X0 X1) \quad (3)$$

Assume the following.

$$v3_membered k1_numbers \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v3_valued_0 X0))) \Rightarrow (m1_subset_1 (k1_seq_1 X0 X1) k1_numbers) \quad (5)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v2_valued_0 X0))) \Rightarrow \\ (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2.(X2 = k11_mesfunc1 X0 \\ X1) \Leftrightarrow (\forall X3.(X3 \in X2) \Leftrightarrow ((X3 \in k9_xtuple_0 X0) \wedge (\neg r1_xreal_0 \\ X1 (k12_supinf_2 X0 X3)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xreal_0 X0) \quad (7)$$

Assume the following.

$$\forall X0.(v3_membered X0) \Rightarrow (v2_membered X0) \quad (8)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.(v3_membered X1) \Rightarrow (\forall X2.(m1_subset_1 \\ X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v3_valued_0 X2)) \end{aligned} \quad (11)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.(v2_membered X1) \Rightarrow (\forall X2.(m1_subset_1 \\ X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v2_valued_0 X2)) \end{aligned} \quad (12)$$

Theorem 1

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\ m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow (\\ \forall X2.(v1_xreal_0 X2) \Rightarrow (\forall X3.(X3 \in k11_mesfunc1 X1 X2) \Leftrightarrow \\ ((X3 \in k1_relset_1 X0 X1) \wedge (\exists X4.(m1_subset_1 X4 k1_numbers) \wedge \\ ((X4 = k1_seq_1 X1 X3) \wedge (\neg r1_xreal_0 X2 X4)))))) \end{aligned}$$