

t3_nat_1 (TMZNPyM- nyPr4D4Gmg4mVTRzHxZf2SuRhnPz)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v3_xxreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (r1_xxreal_0 k6_numbers X0) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 X0 k6_numbers) \wedge (r1_xxreal_0 k6_numbers X0)) \Rightarrow (k6_numbers = k3_xcmplx_0 X0 X1))) \quad (2)$$

Assume the following.

$$\exists X0.(v1_xcmplx_0 X0) \wedge ((v1_xxreal_0 X0) \wedge ((v3_xxreal_0 X0) \wedge (v1_xreal_0 X0))) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X0)) \Rightarrow (X0 = X1))) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (v1_xreal_0 (k3_xcmplx_0 X0 X1)) \quad (5)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_xreal_0 X0) \quad (6)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\neg(k6_numbers \neq X0) \wedge (r1_xxreal_0 X0 k6_numbers))$$