

# t3\_ramsey\_1 (TMNchX- cgrZCGp1mBB9LXgkgHSnzP49AmEDx)

October 27, 2020

Let  $v1\_finset\_1 : \iota \Rightarrow o$  be given. Let  $k2\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. r1\_tarski X0 (k2\_xboole\_0 X0 X1) \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X0 (k1\_zfmisc\_1 X1)) \Leftrightarrow (r1\_tarski X0 X1) \tag{2}$$

Assume the following.

$$\forall X0. (v1\_finset\_1 X0) \Rightarrow (\forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)) \Rightarrow (v1\_finset\_1 X1)) \tag{3}$$

**Theorem 1**

$$\forall X0. \forall X1. \neg(\neg v1\_finset\_1 X0) \wedge (v1\_finset\_1 (k2\_xboole\_0 X0 X1))$$