

t3_rewrite3 (TMYduRCMMsx- Abp6QnbzgQxafCfbYNtVt79V)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (1)$$

Assume the following.

$$\forall X0. (v1_xxreal_0 X0) \Rightarrow (\forall X1. (v1_xxreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X0)) \Rightarrow (X0 = X1)) \quad (2)$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Rightarrow (\forall X1. (v7_ordinal1 X1) \Rightarrow ((X0 \in k2_finseq_1 X1) \Leftrightarrow ((r1_xxreal_0 np_1 X0) \wedge (r1_xxreal_0 X0 X1)))) \quad (3)$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Rightarrow (\forall X1. (v7_ordinal1 X1) \Rightarrow ((\neg r1_xxreal_0 (k1_nat_1 X1 np_1) X0) \Leftrightarrow (r1_xxreal_0 X0 X1))) \quad (4)$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Rightarrow (\forall X1. (v7_ordinal1 X1) \Rightarrow (\forall X2. (v7_ordinal1 X2) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow (r1_xxreal_0 X0 (k2_xcmplx_0 X1 X2)))))) \quad (5)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(\forall X1.(v7_ordinal1\ X1)\Rightarrow(r1_xxreal_0\ X0\ (k2_xcmplx_0\ X0\ X1))) \quad (6)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(\forall X1.(v7_ordinal1\ X1)\Rightarrow(\neg\ (r1_xxreal_0\ X0\ X1)\wedge(\forall X2.(v7_ordinal1\ X2)\Rightarrow(X1\neq k2_xcmplx_0\ X0\ X2)))) \quad (7)$$

Assume the following.

$$((v2_xxreal_0\ np_1)\wedge(m2_subset_1\ np_1\ k1_numbers\ k5_numbers))\wedge\ ((m1_subset_1\ np_1\ k5_numbers)\wedge(m1_subset_1\ np_1\ k1_numbers)) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0\ X0)\wedge((\neg v1_xboole_0\ X1)\wedge\ (m1_subset_1\ X1\ (k1_zfmisc_1\ X0))))\Rightarrow(\forall X2.(m2_subset_1\ X2\ X0\ X1)\Leftrightarrow(m1_subset_1\ X2\ X1)) \quad (9)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (10)$$

Assume the following.

$$\forall X0.((v1_relat_1\ X0)\wedge((v1_funct_1\ X0)\wedge(v1_finseq_1\ X0)))\Rightarrow\ (k4_finseq_1\ X0 = k9_xtuple_0\ X0) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1\ X0)\wedge(m1_subset_1\ X1\ k5_numbers))\Rightarrow\ (k1_nat_1\ X0\ X1 = k2_xcmplx_0\ X0\ X1) \quad (12)$$

Assume the following.

$$(\neg v1_xboole_0\ k4_ordinal1)\wedge(v3_ordinal1\ k4_ordinal1) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1\ X0)\wedge(v7_ordinal1\ X1))\Rightarrow(\ v7_ordinal1\ (k2_xcmplx_0\ X0\ X1)) \quad (14)$$

Assume the following.

$$m1_subset_1\ k5_numbers\ (k1_zfmisc_1\ k1_numbers) \quad (15)$$

Assume the following.

$$\forall X0.((v1_relat_1\ X0)\wedge((v1_funct_1\ X0)\wedge(v1_finseq_1\ X0)))\Rightarrow\ (m1_subset_1\ (k4_finseq_1\ X0)\ (k1_zfmisc_1\ k5_numbers)) \quad (16)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow ((X1 = k3_finseq_1 \\ & X0) \Leftrightarrow (k2_finseq_1 X1 = k9_xtuple_0 X0))) \end{aligned} \quad (17)$$

Assume the following.

$$\forall X0.(v1_relat_1 X0) \Rightarrow ((v1_finseq_1 X0) \Leftrightarrow (\exists X1.(v7_ordinal1 X1) \wedge (k9_xtuple_0 X0 = k2_finseq_1 X1))) \quad (18)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (19)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_xxreal_0 X0) \quad (20)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (v1_xboole_0 X1)) \quad (21)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge ((\\ & v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \Rightarrow ((X0 \in k4_finseq_1 X1) \Rightarrow ((k2_xcmplx_0 \\ & X0 np_1 \in k4_finseq_1 X1) \vee (k3_finseq_1 X1 = X0)))) \end{aligned}$$