

t3_rfunct_1

(TMJq7xV2qrp3iwXDDJ2zhUEd1b4F6KpB27i)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_valued_0 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k4_rfunct_1 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k5_xcmplx_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. k6_subset_1 X0 X1 = k4_xboole_0 X0 X1 \quad (1)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 X0))) \Rightarrow ((v1_relat_1 (k4_rfunct_1 X0)) \wedge (v1_funct_1 (k4_rfunct_1 X0))) \quad (2)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. (X2 = k8_relat_1 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in k9_xtuple_0 X0) \wedge (k1_funct_1 X0 X3 \in X1)))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k4_xboole_0 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in X0) \wedge (\neg X3 \in X1))) \quad (4)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 X0))) \Rightarrow (\forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((X1 = k4_rfunct_1 X0) \Leftrightarrow ((k9_xtuple_0 X1 = k6_subset_1 (k9_xtuple_0 X0) (k8_relat_1 X0 (k1_tarski k6_numbers))) \wedge (\forall X2. (X2 \in k9_xtuple_0 X1) \Rightarrow (k1_funct_1 X1 X2 = k5_xcmplx_0 (k1_funct_1 X0 X2)))))) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (6)$$

Theorem 1

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 X0) \Rightarrow \\ & (\forall X2. ((v1_relat_1 X2) \wedge ((v1_funct_1 X2) \wedge (v1_valued_0 \\ & X2)))) \Rightarrow (\neg (X1 \in k9_xtuple_0 (k4_rfunct_1 X2)) \wedge (k1_funct_1 X2 X1 = \\ & k6_numbers)))) \end{aligned}$$