

t3_roughs_1 (TMQAf-
SLrya7n4gYNbHMzpzDwVoYa3H95MsZ)

October 27, 2020

Let $v1_relat_2 : \iota \Rightarrow o$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_partfun1 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_relat_1 : \iota \Rightarrow \iota$ be given. Let $k1_relat_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_relat_1 X0) \Rightarrow ((v1_relat_2 X0) \Leftrightarrow (r1_tarski (k4_relat_1 (k1_relat_1 X0)) X0)) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow ((r1_relset_1 X0 X1 X2 X3) \Leftrightarrow (r1_tarski X2 X3)) \quad (2)$$

Assume the following.

$$\forall X0.k6_partfun1 X0 = k4_relat_1 X0 \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1_partfun1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 X0)))) \Rightarrow (k1_relat_1 X1 = X0) \quad (4)$$

Assume the following.

$$\forall X0.(v1_partfun1 (k6_partfun1 X0) X0) \wedge (m1_subset_1 (k6_partfun1 X0) (k1_zfmisc_1 (k2_zfmisc_1 X0 X0))) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \quad (6)$$

Theorem 1

$$\forall X0. \forall X1. ((v1_relat_2 X1) \wedge ((v1_partfun1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 X0)))))) \Rightarrow (r1_relset_1 X0 X0 (k6_partfun1 X0) X1)$$