

t3_tdlat_3

(TMUZz1F4nVVXYrhrXu16asXhqYmhzspgQq2)

October 27, 2020

Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k1_tops_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_pre_topc : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (k3_subset_1 X0 (k3_subset_1 X0 X1) = X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (m1_subset_1 (k3_subset_1 X0 X1) (k1_zfmisc_1 X0)) \quad (2)$$

Assume the following.

$$\forall X0. (l1_pre_topc X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (k1_tops_1 X0 X1 = k3_subset_1 (u1_struct_0 X0) X0) (k2_pre_topc X0 (k3_subset_1 (u1_struct_0 X0) X1)))) \quad (3)$$

Theorem 1

$$\forall X0. ((v2_pre_topc X0) \wedge (l1_pre_topc X0)) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (k1_tops_1 X0 (k3_subset_1 (u1_struct_0 X0) X1) = k3_subset_1 (u1_struct_0 X0) (k2_pre_topc X0 X1)))$$