

t3_topreal4

(TMcFn1oZx2ZdtZZtn2arGfWccviWM9RpBk1)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $r1_topreal4 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_topreal1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v6_membered : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v2_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rlvect_1 : \iota \Rightarrow o$ be given. Let $v6_rlvect_1 : \iota \Rightarrow o$ be given. Let $v7_rlvect_1 : \iota \Rightarrow o$ be given. Let $v8_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rltopsp1 : \iota \Rightarrow o$ be given. Let $l1_rltopsp1 : \iota \Rightarrow o$ be given. Let $l1_rlvect_1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid \\ np_2)))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid \\ np_2))) \Rightarrow (\forall X2.(m1_subset_1 X2 (u1_struct_0 (k15_euclid \\ np_2))) \Rightarrow ((r1_topreal4 X0 X1 X2) \Rightarrow (r1_topreal1 (k15_euclid np_2) \\ X1 X2 X0)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.((v2_pre_topc X0) \wedge (l1_pre_topc X0)) \Rightarrow (\forall X1. \\ (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (\forall X2. \\ (m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (\forall X3.(m1_subset_1 X3 \\ (u1_struct_0 X0)) \Rightarrow ((r1_topreal1 X0 X2 X3 X1) \Rightarrow ((X2 \in X1) \wedge (X3 \in X1))))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} ((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge \\ ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (4)$$

Assume the following.

$$v6_membered\ k4_ordinal1 \tag{5}$$

Assume the following.

$$\begin{aligned} \forall X0.(v7_ordinal1\ X0) \Rightarrow & ((v2_pre_topc\ (k15_euclid\ X0)) \wedge \\ & ((v13_algstr_0\ (k15_euclid\ X0)) \wedge ((v2_rlvect_1\ (k15_euclid\ X0)) \wedge \\ & ((v3_rlvect_1\ (k15_euclid\ X0)) \wedge ((v4_rlvect_1\ (k15_euclid\ X0)) \wedge \\ & ((v5_rlvect_1\ (k15_euclid\ X0)) \wedge ((v6_rlvect_1\ (k15_euclid\ X0)) \wedge \\ & ((v7_rlvect_1\ (k15_euclid\ X0)) \wedge ((v8_rlvect_1\ (k15_euclid\ X0)) \wedge \\ & (v5_rltopsp1\ (k15_euclid\ X0)))))))))) \end{aligned} \tag{6}$$

Assume the following.

$$\forall X0.(l1_rltopsp1\ X0) \Rightarrow ((l1_rlvect_1\ X0) \wedge (l1_pre_topc\ X0)) \tag{7}$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0) \Rightarrow ((v5_rltopsp1\ (k15_euclid\ X0)) \wedge (l1_rltopsp1\ (k15_euclid\ X0))) \tag{8}$$

Assume the following.

$$\forall X0.(v6_membered\ X0) \Rightarrow (\forall X1.(m1_subset_1\ X1\ X0) \Rightarrow (v7_ordinal1\ X1)) \tag{9}$$

Theorem 1

$$\begin{aligned} \forall X0.(m1_subset_1\ X0\ (k1_zfmisc_1\ (u1_struct_0\ (k15_euclid \\ np_2)))) \Rightarrow & (\forall X1.(m1_subset_1\ X1\ (u1_struct_0\ (k15_euclid \\ np_2)))) \Rightarrow & (\forall X2.(m1_subset_1\ X2\ (u1_struct_0\ (k15_euclid \\ np_2)))) \Rightarrow & ((r1_topreal4\ X0\ X1\ X2) \Rightarrow ((X1 \in X0) \wedge (X2 \in X0))) \end{aligned}$$