

t3_toprns_1
(TMGuwRgBmkPJjiQxfF9WkfW6z2q6x4fanGE)

October 27, 2020

Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_toprns_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k8_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_struct_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m2_subset_1 X0 k1_numbers k5_numbers) \Rightarrow (\forall X1. \\ & ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 k5_numbers (u1_struct_0 (k15_euclid \\ & X0))) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers (\\ & u1_struct_0 (k15_euclid X0))))))) \Rightarrow ((v1_toprns_1 X1 X0) \Leftrightarrow (\forall X2. \\ & \neg (X2 \in k5_numbers) \wedge (k1_funct_1 X1 X2 = k4_struct_0 (k15_euclid \\ & X0)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2. (m2_subset_1 \\ & X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \tag{4}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (((v1_funct_1 X1) \wedge ((v1_funct_2 \\ & X1 k5_numbers X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers \\ & X0)))))) \wedge (v7_ordinal1 X2) \Rightarrow (k8_nat_1 X0 X1 X2 = k1_funct_1 X1 X2) \end{aligned} \tag{5}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (6)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (7)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (8)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (9)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (10)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m2_subset_1 X0 k1_numbers k5_numbers) \Rightarrow (\forall X1. \\ & ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 k5_numbers (u1_struct_0 (k15_euclid \\ & X0))) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers (\\ & u1_struct_0 (k15_euclid X0))))))) \Rightarrow ((v1_toprns_1 X1 X0) \Leftrightarrow (\forall X2. \\ & (m2_subset_1 X2 k1_numbers k5_numbers) \Rightarrow (k8_nat_1 (u1_struct_0 \\ & (k15_euclid X0)) X1 X2 \neq k4_struct_0 (k15_euclid X0)))))) \end{aligned}$$