

t3_waybel15
(TMKKLXZrrW6ju5CB5F87gkkuCVL5GtskDHS)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v5_orders_2 : \iota \Rightarrow o$ be given. Let $v2_yellow_0 : \iota \Rightarrow o$ be given. Let $l1_orders_2 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $r1_orders_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_yellow_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.((\neg v2_struct_0 X0) \wedge ((v5_orders_2 X0) \wedge ((v2_yellow_0 X0) \wedge (l1_orders_2 X0)))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (r1_orders_2 X0 X1 (k4_yellow_0 X0))) \quad (1)$$

Assume the following.

$$\forall X0.(l1_orders_2 X0) \Rightarrow (m1_subset_1 (k4_yellow_0 X0) (u1_struct_0 X0)) \quad (2)$$

Assume the following.

$$\forall X0.(l1_orders_2 X0) \Rightarrow ((v5_orders_2 X0) \Leftrightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow ((r1_orders_2 X0 X1 X2) \wedge (r1_orders_2 X0 X2 X1)) \Rightarrow (X1 = X2)))))) \quad (3)$$

Theorem 1

$$\forall X0.((\neg v2_struct_0 X0) \wedge ((v5_orders_2 X0) \wedge ((v2_yellow_0 X0) \wedge (l1_orders_2 X0)))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow ((r1_orders_2 X0 (k4_yellow_0 X0) X1) \Rightarrow (X1 = k4_yellow_0 X0)))$$