

t402\_xxreal\_1

(TMZbhRToDvUvz4amggchf79F57iasxZj5jQ)

October 27, 2020

Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_xxreal\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_xxreal\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $k1\_xxreal\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xxreal\_0 X2) \Rightarrow ((\neg r1\_xxreal\_0 X1 X0) \Rightarrow (k6\_subset\_1 (k3\_xxreal\_1 \\ & X0 X2) (k4\_xxreal\_1 X0 X1) = k1\_xxreal\_1 X1 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (k1\_xxreal\_1 X0 X0 = k1\_tarski X0) \quad (2)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (( \\ & \neg r1\_xxreal\_0 X1 X0) \Rightarrow (k6\_subset\_1 (k3\_xxreal\_1 X0 X1) (k4\_xxreal\_1 \\ & X0 X1) = k1\_tarski X1))) \end{aligned}$$