

t40_diff_2

(TMMU3D5mXF5ZKTdeKcFjDDF6WuMd619GGhL)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k9_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_diff_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_real_1 : \iota \Rightarrow \iota$ be given. Let $k8_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $np_0 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k3_xcmplx_0 np_1 X0 = X0) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k3_xcmplx_0 X0 k6_numbers = k6_numbers) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0) \wedge (v1_xcmplx_0 X1)) \Rightarrow (k2_xcmplx_0 (k4_xcmplx_0 X0) (k4_xcmplx_0 X1) = k4_xcmplx_0 (k2_xcmplx_0 X0 X1)) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1_xcmplx_0 X0) \wedge ((v1_xcmplx_0 X1) \wedge (v1_xcmplx_0 X2))) \Rightarrow (k3_xcmplx_0 (k2_xcmplx_0 X0 X1) X2 = k2_xcmplx_0 (k3_xcmplx_0 X0 X2) (k3_xcmplx_0 X1 X2)) \quad (5)$$

Assume the following.

$$\begin{aligned} & ((v2_xreal_0 \ np_2) \wedge (m2_subset_1 \ np_2 \ k1_numbers \ k5_numbers)) \wedge \\ & ((m1_subset_1 \ np_2 \ k5_numbers) \wedge (m1_subset_1 \ np_2 \ k1_numbers)) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0. (v1_xcmplx_0 \ X0) \Rightarrow (k3_xcmplx_0 \ X0 \ (k4_xcmplx_0 \ np_1) = k4_xcmplx_0 \ X0) \quad (7)$$

Assume the following.

$$\begin{aligned} & ((v2_xreal_0 \ np_1) \wedge (m2_subset_1 \ np_1 \ k1_numbers \ k5_numbers)) \wedge \\ & ((m1_subset_1 \ np_1 \ k5_numbers) \wedge (m1_subset_1 \ np_1 \ k1_numbers)) \end{aligned} \quad (8)$$

Assume the following.

$$(m2_subset_1 \ np_0 \ k1_numbers \ k5_numbers) \wedge ((m1_subset_1 \ np_0 \ k5_numbers) \wedge (m1_subset_1 \ np_0 \ k1_numbers)) \quad (9)$$

Assume the following.

$$v1_xboole_0 \ np_0 \quad (10)$$

Assume the following.

$$k4_xcmplx_0 \ (k4_xcmplx_0 \ np_1) = np_1 \quad (11)$$

Assume the following.

$$k4_xcmplx_0 \ np_0 = np_0 \quad (12)$$

Assume the following.

$$k2_xcmplx_0 \ np_1 \ (k4_xcmplx_0 \ np_1) = np_0 \quad (13)$$

Assume the following.

$$\forall X0. \forall X1. ((m1_subset_1 \ X0 \ k1_numbers) \wedge (v1_xreal_0 \ X1)) \Rightarrow (k8_real_1 \ X0 \ X1 = k3_xcmplx_0 \ X0 \ X1) \quad (14)$$

Assume the following.

$$\forall X0. \forall X1. ((m1_subset_1 \ X0 \ k1_numbers) \wedge (v1_xreal_0 \ X1)) \Rightarrow (k7_real_1 \ X0 \ X1 = k2_xcmplx_0 \ X0 \ X1) \quad (15)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (16)$$

Assume the following.

$$\forall X0. (m1_subset_1 \ X0 \ k1_numbers) \Rightarrow (k1_real_1 \ X0 = k4_xcmplx_0 \ X0) \quad (17)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers)\wedge(v1_xreal_0 X1))\Rightarrow(m1_subset_1 (k8_real_1 X0 X1) k1_numbers) \quad (18)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(m1_subset_1 (k1_real_1 X0) k1_numbers) \quad (19)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers)\wedge(v1_xreal_0 X1))\Rightarrow(m1_subset_1 (k10_real_1 X0 X1) k1_numbers) \quad (20)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0)\wedge(v1_xcmplx_0 X1))\Rightarrow(k3_xcmplx_0 X0 X1 = k3_xcmplx_0 X1 X0) \quad (21)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(v1_xcmplx_0 X0) \quad (22)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (23)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(\forall X1.(m1_subset_1 \\ & X1 k1_numbers)\Rightarrow(\forall X2.((v1_funct_1 X2)\wedge((v1_funct_2 X2 \\ & k1_numbers k1_numbers)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ & k1_numbers k1_numbers))))\Rightarrow((\forall X3.(m1_subset_1 X3 k1_numbers)\Rightarrow \\ & ((k3_funct_2 k1_numbers k1_numbers X2 X3 = k10_real_1 X0 X3)\wedge((\\ & k7_real_1 X3 (k10_real_1 X1 np_2)\neq k6_numbers)\wedge(k9_real_1 X3 \\ & (k10_real_1 X1 np_2)\neq k6_numbers))))\Rightarrow(\forall X3.(m1_subset_1 \\ & X3 k1_numbers)\Rightarrow(k1_funct_1 (k5_diff_1 X2 X1) X3 = k10_real_1 (k1_real_1 \\ & (k8_real_1 X0 X1) (k8_real_1 (k9_real_1 X3 (k10_real_1 X1 np_2)) \\ & (k7_real_1 X3 (k10_real_1 X1 np_2)))))))))) \end{aligned}$$