

t40_euclid_3 (TMZm-
FkX1sSTnyLN11RoDUH5emqT5BiFi5Vn)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_numbers : \iota$ be given. Let $k4_complex1 : \iota \Rightarrow \iota$ be given. Let $k1_complex2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_real_1 : \iota \Rightarrow \iota$ be given. Let $k8_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_complex1 : \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_complex1 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_subset_1 X0 k2_numbers) \Rightarrow (\forall X1.(m1_subset_1 \\ X1 k2_numbers) \Rightarrow (k1_complex2 X0 X1 = k2_xcmplx_0 (k7_real_1 (k8_real_1 \\ (k3_complex1 X0) (k3_complex1 X1)) (k8_real_1 (k4_complex1 X0) \\ (k4_complex1 X1))) (k3_xcmplx_0 (k7_real_1 (k1_real_1 (k8_real_1 \\ (k3_complex1 X0) (k4_complex1 X1)))) (k8_real_1 (k4_complex1 X0) \\ (k3_complex1 X1))) k7_complex1))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((k3_complex1 \\ (k2_xcmplx_0 X0 (k3_xcmplx_0 X1 k7_complex1)) = X0) \wedge (k4_complex1 \\ (k2_xcmplx_0 X0 (k3_xcmplx_0 X1 k7_complex1)) = X1))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers) \wedge (v1_xreal_0 X1)) \Rightarrow (m1_subset_1 (k8_real_1 X0 X1) k1_numbers) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers) \wedge (v1_xreal_0 X1)) \Rightarrow (m1_subset_1 (k7_real_1 X0 X1) k1_numbers) \quad (4)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (m1_subset_1 (k4_complex1 X0) k1_numbers) \quad (5)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (m1_subset_1 (k3_complex1 X0) k1_numbers) \quad (6)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (m1_subset_1 (k1_real_1 X0) k1_numbers) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers) \wedge (v1_xreal_0 X1)) \Rightarrow (k8_real_1 X0 X1 = k8_real_1 X1 X0) \quad (8)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k2_numbers) \Rightarrow (v1_xcmplx_0 X0) \quad (9)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (10)$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 k2_numbers) \Rightarrow (\forall X1.(m1_subset_1 X1 k2_numbers) \Rightarrow (k4_complex1 (k1_complex2 X0 X1) = k7_real_1 (k1_real_1 (k8_real_1 (k3_complex1 X0) (k4_complex1 X1))) (k8_real_1 (k4_complex1 X0) (k3_complex1 X1))))$$