

t40_facirc_1 (TMPy-
PaT7qLwrd5FEEqzokERgeDJAXBKSMP6)

October 27, 2020

Let $k2_msafree2 : \iota \Rightarrow \iota$ be given. Let $k5_circcomb : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k3_msafree2 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \Rightarrow ((k2_msafree2 (k5_circcomb X0 X1) = k10_xtuple_0 X1) \wedge (k3_msafree2 (k5_circcomb X0 X1) = k1_tarski (k4_tarski X1 X0))) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. k10_xtuple_0 (k10_finseq_1 X0 X1) = k2_tarski X0 X1 \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 (k10_finseq_1 X0 X1)) \wedge (v1_funct_1 (k10_finseq_1 X0 X1)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. v1_finseq_1 (k10_finseq_1 X0 X1) \quad (4)$$

Theorem 1

$$\forall X0. \forall X1. \forall X2. k2_msafree2 (k5_circcomb X0 (k10_finseq_1 X1 X2)) = k2_tarski X1 X2$$