

t40_lpspace1

(TMHEXr7fWkygLwhUPrMazFb2izV961zgQoy)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_numbers : \iota$ be given. Let $v10_valued_0 : \iota \Rightarrow o$ be given. Let $v6_supinf_2 : \iota \Rightarrow o$ be given. Let $v4_measure1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $k9_lpspace1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k13_lpspace1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_lpspace1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned}
& \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((\neg v1_xboole_0 X1) \wedge \\
& ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\
& (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2. ((v1_funct_1 X2) \wedge ((v1_funct_2 \\
& X2 X1 k7_numbers) \wedge ((v10_valued_0 X2) \wedge ((v6_supinf_2 X2) \wedge ((v4_measure1 \\
& X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers)))))) \Rightarrow \\
& (\forall X3. ((v1_funct_1 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\
& X0 k1_numbers)))) \Rightarrow (\forall X4. ((v1_funct_1 X4) \wedge (m1_subset_1 \\
& X4 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow (((X3 \in k9_lpspace1 \\
& X0 X1 X2) \wedge (X4 \in k9_lpspace1 X0 X1 X2)) \Rightarrow ((k13_lpspace1 X0 X1 X2 X3 = \\
& k13_lpspace1 X0 X1 X2 X4) \Leftrightarrow (r1_lpspace1 X0 X1 X2 X3 X4))))))
\end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned}
& \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((\neg v1_xboole_0 X1) \wedge \\
& ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\
& (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2. ((v1_funct_1 X2) \wedge ((v1_funct_2 \\
& X2 X1 k7_numbers) \wedge ((v10_valued_0 X2) \wedge ((v6_supinf_2 X2) \wedge ((v4_measure1 \\
& X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers)))))) \Rightarrow \\
& (\forall X3. ((v1_funct_1 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\
& X0 k1_numbers)))) \Rightarrow (\forall X4. ((v1_funct_1 X4) \wedge (m1_subset_1 \\
& X4 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow (((X3 \in k9_lpspace1 \\
& X0 X1 X2) \wedge (X4 \in k9_lpspace1 X0 X1 X2)) \Rightarrow ((r1_lpspace1 X0 X1 X2 X4 X3) \Leftrightarrow \\
& (X4 \in k13_lpspace1 X0 X1 X2 X3))))))
\end{aligned} \tag{2}$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((\neg v1_xboole_0 X1) \wedge \\ & ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge ((v1_funct_2 \\ & X2 X1 k7_numbers) \wedge ((v10_valued_0 X2) \wedge ((v6_supinf_2 X2) \wedge ((v4_measure1 \\ & X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers)))))) \Rightarrow \\ & (\forall X3.((v1_funct_1 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 k1_numbers)))) \Rightarrow (\forall X4.((v1_funct_1 X4) \wedge (m1_subset_1 \\ & X4 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow (((X3 \in k9_lpspace1 \\ & X0 X1 X2) \wedge (X4 \in k9_lpspace1 X0 X1 X2)) \Rightarrow ((k13_lpspace1 X0 X1 X2 X3 = \\ & k13_lpspace1 X0 X1 X2 X4) \Leftrightarrow (X4 \in k13_lpspace1 X0 X1 X2 X3)))))) \end{aligned}$$