

t40_partfun1 (TMT- cut6fkWwDw15DMLetWJnNXp3Hiab9Qxu)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 \\ & X2)) \Rightarrow ((r1_tarski (k1_relset_1 X0 (k3_partfun1 X2 X0 X1)) (k9_xtuple_0 \\ & X2)) \wedge (r1_tarski (k10_xtuple_0 (k3_partfun1 X2 X0 X1)) (k10_xtuple_0 \\ & X2))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((v1_relat_1 X0) \wedge (v1_funct_1 \\ & X0)) \Rightarrow ((v1_funct_1 (k3_partfun1 X0 X1 X2)) \wedge (m1_subset_1 (k3_partfun1 \\ & X0 X1 X2) (k1_zfmisc_1 (k2_zfmisc_1 X1 X2)))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (\\ & (v1_partfun1 X1 X0) \Leftrightarrow (k1_relset_1 X0 X1 = X0)) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \end{aligned} \tag{4}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \end{aligned} \tag{5}$$

Theorem 1

$$\forall X0.\forall X1.\forall X2.((v1_relat_1 X2)\wedge(v1_funct_1 X2))\Rightarrow((v1_partfun1 (k3_partfun1 X2 X0 X1) X0)\Rightarrow(r1_tarski X0 (k9_xtuple_0 X2)))$$