

t40_setfam_1
(TMa3rEq6ckThaKnQR2e4tqn5nBa4bcmZ7bP)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k7_setfam_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k3_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_subset_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 (k1_zfmisc_1 X1)) \Leftrightarrow (r1_tarski X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. k4_xboole_0 X0 k1_xboole_0 = X0 \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski (k1_tarski X0) X1) \Leftrightarrow (X0 \in X1) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. r1_tarski X0 X0 \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (k3_subset_1 X0 (k3_subset_1 X0 X1) = X1) \quad (6)$$

Assume the following.

$$\forall X0. \neg v1_xboole_0 (k1_zfmisc_1 X0) \quad (7)$$

Assume the following.

$$\forall X0. m1_subset_1 (k1_subset_1 X0) (k1_zfmisc_1 X0) \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 \\ & X0))) \Rightarrow (\forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k1_zfmisc_1 \\ & X0))) \Rightarrow ((X2 = k7_setfam_1 X0 X1) \Leftrightarrow (\forall X3. (m1_subset_1 X3 (k1_zfmisc_1 \\ & X0)) \Rightarrow ((X3 \in X2) \Leftrightarrow (k3_subset_1 X0 X3 \in X1)))))) \end{aligned} \quad (9)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (k3_subset_1 X0 X1 = k4_xboole_0 X0 X1) \quad (10)$$

Assume the following.

$$k1_xboole_0 = the (\lambda X0 : \iota.v1_xboole_0 X0) \quad (11)$$

Assume the following.

$$\forall X0. k1_subset_1 X0 = k1_xboole_0 \quad (12)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (13)$$

Theorem 1

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0))) \Rightarrow ((X1 = k1_tarski X0) \Rightarrow (k7_setfam_1 X0 X1 = k1_tarski k1_xboole_0))$$