

t40\_tex\_2

(TMYbxsuPkXco27xnryqPjSRRcfqN1xXA6hf)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v2\_pre\_topc : \iota \Rightarrow o$  be given. Let  $l1\_pre\_topc : \iota \Rightarrow o$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $v3\_tex\_2 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $v2\_tex\_2 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_domain\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $l1\_struct\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (X0 = k1\_xboole\_0) \quad (1)$$

Assume the following.

$$\forall X0.((\neg v2\_struct\_0 X0) \wedge ((v2\_pre\_topc X0) \wedge (l1\_pre\_topc X0))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 X0)) \Rightarrow (v2\_tex\_2 (k6\_domain\_1 (u1\_struct\_0 X0) X1) X0)) \quad (2)$$

Assume the following.

$$\forall X0.r1\_tarski k1\_xboole\_0 X0 \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0 X0) \wedge (m1\_subset\_1 X1 X0)) \Rightarrow (k6\_domain\_1 X0 X1 = k1\_tarski X1) \quad (4)$$

Assume the following.

$$\forall X0.\neg v1\_xboole\_0 (k1\_tarski X0) \quad (5)$$

Assume the following.

$$\forall X0.((\neg v2\_struct\_0 X0) \wedge (l1\_struct\_0 X0)) \Rightarrow (\neg v1\_xboole\_0 (u1\_struct\_0 X0)) \quad (6)$$

Assume the following.

$$\forall X0.\exists X1.m1\_subset\_1 X1 X0 \quad (7)$$

Assume the following.

$$\forall X0.(l1\_pre\_topc\ X0)\Rightarrow(l1\_struct\_0\ X0) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0\ X0)\wedge(m1\_subset\_1\ X1\ X0))\Rightarrow(m1\_subset\_1\ (k6\_domain\_1\ X0\ X1)\ (k1\_zfmisc\_1\ X0)) \quad (9)$$

Assume the following.

$$\begin{aligned} \forall X0.(l1\_pre\_topc\ X0)\Rightarrow(\forall X1.(m1\_subset\_1\ X1\ (k1\_zfmisc\_1 \\ (u1\_struct\_0\ X0)))\Rightarrow((v3\_tex\_2\ X1\ X0)\Leftrightarrow((v2\_tex\_2\ X1\ X0)\wedge(\forall X2. \\ (m1\_subset\_1\ X2\ (k1\_zfmisc\_1\ (u1\_struct\_0\ X0)))\Rightarrow((v2\_tex\_2 \\ X2\ X0)\wedge(r1\_tarski\ X1\ X2))\Rightarrow(X1 = X2)))))) \quad (10) \end{aligned}$$

**Theorem 1**

$$\forall X0.((\neg v2\_struct\_0\ X0)\wedge((v2\_pre\_topc\ X0)\wedge(l1\_pre\_topc\ X0)))\Rightarrow(\forall X1.((v1\_xboole\_0\ X1)\wedge(m1\_subset\_1\ X1\ (k1\_zfmisc\_1\ (u1\_struct\_0\ X0))))\Rightarrow(\neg v3\_tex\_2\ X1\ X0))$$