

t40_xxreal_3
(TMKT12fCtReBtFZHcdJCsXNsyj3VgDi24E5)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k3_xxreal_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xxreal_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xxreal_3 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (k1_xxreal_3 X0 (k2_xxreal_3 X0) = k6_numbers) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (\forall X3.(v1_xxreal_0 X3) \Rightarrow (((r1_xxreal_0 \\ & X0 X1) \wedge (r1_xxreal_0 X2 X3)) \Rightarrow (r1_xxreal_0 (k1_xxreal_3 X0 X2) (\\ & k1_xxreal_3 X1 X3)))))) \quad (2) \end{aligned}$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (r1_xxreal_0 X0 X0) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (v1_xxreal_0 (k2_xxreal_3 X0)) \quad (4)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (k3_xxreal_3 X0 X1 = k1_xxreal_3 X0 (k2_xxreal_3 X1))) \quad (5)$$

Theorem 1

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow (r1_xxreal_0 k6_numbers (k3_xxreal_3 X1 X0))))$$