

t411\_xxreal\_1  
(TMVCY1Dm8DVZgxhmcerscsnQn4ikzgvhZVJ)

October 27, 2020

Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $k6\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xxreal\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xxreal\_0 : \iota$  be given. Let  $k3\_xxreal\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (k6\_subset\_1 \\ (k1\_xxreal\_1 X0 k1\_xxreal\_0) (k3\_xxreal\_1 X1 k1\_xxreal\_0) = k1\_xxreal\_1 \\ X0 X1)) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (k1\_xxreal\_1 X0 X0 = k1\_tarski X0) \tag{2}$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (v1\_xxreal\_0 X0) \tag{3}$$

**Theorem 1**

$$\begin{aligned} \forall X0.(v1\_xreal\_0 X0) \Rightarrow (k6\_subset\_1 (k1\_xxreal\_1 X0 k1\_xxreal\_0) \\ (k3\_xxreal\_1 X0 k1\_xxreal\_0) = k1\_tarski X0) \end{aligned}$$