

t412_xxreal_1

(TMZb8YcYBPqLPdAWu6NbWyMtyMFAx55uTUL)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xxreal_0 : \iota$ be given. Let $k4_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k1_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (k6_subset_1 \\ (k2_xxreal_1 X0 k1_xxreal_0) (k4_xxreal_1 X1 k1_xxreal_0) = k1_xxreal_1 \\ X0 X1)) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (k1_xxreal_1 X0 X0 = k1_tarski X0) \tag{2}$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xxreal_0 X0) \tag{3}$$

Theorem 1

$$\begin{aligned} \forall X0.(v1_xreal_0 X0) \Rightarrow (k6_subset_1 (k2_xxreal_1 X0 k1_xxreal_0) \\ (k4_xxreal_1 X0 k1_xxreal_0) = k1_tarski X0) \end{aligned}$$