

t417_xxreal_1
 (TMQsfu4GCxydQqr3Mu8h4yHwpeaF2CWpmYV)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (k1_xxreal_1 X0 X0 = k1_tarski X0) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (\forall X3.(v1_xxreal_0 X3) \Rightarrow ((r1_xxreal_0 \\ & X2 X3) \Rightarrow ((r1_xxreal_0 X1 X0) \vee (k3_xboole_0 (k3_xxreal_1 X0 X2) (\\ & k1_xxreal_1 X1 X3) = k1_xxreal_1 X1 X2)))))) \quad (2) \end{aligned}$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow ((r1_xxreal_0 X1 X2) \Rightarrow ((r1_xxreal_0 X1 X0) \vee (\\ & k3_xboole_0 (k3_xxreal_1 X0 X1) (k1_xxreal_1 X1 X2) = k1_tarski \\ & X1)))))) \end{aligned}$$