

t41_borsuk_5

(TMLovn2xwgDrcavZe1qWqeREDKFnAKmYMfz)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k3_topmetr : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k4_rcomp_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xxreal_0 : \iota$ be given. Let $v4_pre_topc : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $v2_rcomp_1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_subset_1 X0 (k1_zfmisc_1 k1_numbers)) \Rightarrow (\forall X1. \\ (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 k3_topmetr))) \Rightarrow ((X0 = \\ X1) \Rightarrow ((v2_rcomp_1 X0) \Leftrightarrow (v4_pre_topc X1 k3_topmetr)))) \end{aligned} \quad (1)$$

Assume the following.

$$u1_struct_0 k3_topmetr = k1_numbers \quad (2)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v2_rcomp_1 (k4_rcomp_1 k2_xxreal_0 X0)) \quad (3)$$

Theorem 1

$$\begin{aligned} \forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 k3_topmetr))) \Rightarrow \\ (\forall X1.(v1_xreal_0 X1) \Rightarrow ((X0 = k4_rcomp_1 k2_xxreal_0 X1) \Rightarrow \\ (v4_pre_topc X0 k3_topmetr))) \end{aligned}$$