

t41_cqc_the3 (TM- SNXQb3vpamj5skrsEamoW1eVm2Vw6fH2t)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k3_cqc_lang : \iota \Rightarrow \iota$ be given. Let $v6_qc_lang1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_cqc_the3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v2_cqc_the1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k8_cqc_lang : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $r3_cqc_the1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_qc_lang2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_subset_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 \\ (k3_cqc_lang X0))) \Rightarrow (\forall X2.(m2_subset_1 X2 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (\forall X3.(m2_subset_1 X3 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (((v6_qc_lang1 X2 X0) \wedge (r3_cqc_the1 X0 (\\ k4_subset_1 (k3_cqc_lang X0) X1 (k6_domain_1 (k3_cqc_lang X0) \\ X2)) X3)) \Rightarrow (r3_cqc_the1 X0 X1 (k8_cqc_lang X0 X2 X3)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (2)$$

Assume the following.

$$\forall X0.k2_xboole_0 X0 k1_xboole_0 = X0 \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.((m1_qc_lang1 X0) \wedge ((m1_subset_1 \\ X1 (k3_cqc_lang X0)) \wedge (m1_subset_1 X2 (k3_cqc_lang X0)))) \Rightarrow (k8_cqc_lang \\ X0 X1 X2 = k2_qc_lang2 X0 X1 X2) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1_subset_1 X1 (k1_zfmisc_1 X0))\wedge(m1_subset_1 X2 (k1_zfmisc_1 X0)))\Rightarrow(k4_subset_1 X0 X1 X2 = k2_xboole_0 X1 X2) \quad (6)$$

Assume the following.

$$\forall X0.\exists X1.(m1_subset_1 X1 (k1_zfmisc_1 X0))\wedge(v1_xboole_0 X1) \quad (7)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0)\Rightarrow(\neg v1_xboole_0 (k3_cqc_lang X0)) \quad (8)$$

Assume the following.

$$\forall X0.v1_xboole_0 (k1_subset_1 X0) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((\neg v1_xboole_0 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 X0))))\Rightarrow(\forall X2.(m2_subset_1 X2 X0 X1)\Rightarrow(m1_subset_1 X2 X0)) \quad (10)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0)\Rightarrow(\neg v1_xboole_0 (k9_qc_lang1 X0)) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge(m1_subset_1 X1 X0))\Rightarrow(m1_subset_1 (k6_domain_1 X0 X1) (k1_zfmisc_1 X0)) \quad (12)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0)\Rightarrow(m1_subset_1 (k3_cqc_lang X0) (k1_zfmisc_1 (k9_qc_lang1 X0))) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1_qc_lang1 X0)\wedge((m1_subset_1 X1 (k9_qc_lang1 X0))\wedge(m1_subset_1 X2 (k9_qc_lang1 X0))))\Rightarrow(m1_subset_1 (k2_qc_lang2 X0 X1 X2) (k9_qc_lang1 X0)) \quad (14)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 (k9_qc_lang1 X0))\Rightarrow((v2_cqc_the1 X1 X0)\Leftrightarrow(r3_cqc_the1 X0 (k1_subset_1 (k3_cqc_lang X0) X1)))) \quad (15)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m2_subset_1\ X1\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (\forall X2.(m2_subset_1\ X2\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow ((r1_cqc_the3\ X0\ X1\ X2) \Leftrightarrow (r3_cqc_the1\ X0 \\ (k6_domain_1\ (k3_cqc_lang\ X0)\ X1\ X2)))) \end{aligned} \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.k2_xboole_0\ X0\ X1 = k2_xboole_0\ X1\ X0 \quad (17)$$

Theorem 1

$$\begin{aligned} \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m2_subset_1\ X1\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (\forall X2.(m2_subset_1\ X2\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (((v6_qc_lang1\ X1\ X0) \wedge (r1_cqc_the3\ X0\ X1 \\ X2)) \Rightarrow (v2_cqc_the1\ (k8_cqc_lang\ X0\ X1\ X2)\ X0)))) \end{aligned}$$