

t41_mesfun7c

(TMJY9o2sDBpRknyHbHhypBrHMiSjXVPy6Ye)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v6_supinf_2 : \iota \Rightarrow o$ be given. Let $k47_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\ & m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow (\\ & \forall X2.((v1_funct_1 X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 k1_numbers)))) \Rightarrow ((\forall X3.(X3 \in k9_subset_1 X0 (k1_relset_1 \\ & X0 X1) (k1_relset_1 X0 X2)) \Rightarrow (r1_xxreal_0 (k1_seq_1 X2 X3) (k1_seq_1 \\ & X1 X3))) \Rightarrow (v6_supinf_2 (k47_valued_1 X0 k1_numbers k1_numbers \\ & X1 X2)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow (k9_subset_1 X0 X1 X2 = k3_xboole_0 X1 X2) \tag{2}$$

Assume the following.

$$\forall X0.\forall X1.v1_relat_1 (k2_zfmisc_1 X0 X1) \tag{3}$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (m1_subset_1 (k1_relset_1 X0 X1) (k1_zfmisc_1 X0)) \tag{4}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(X2 = k3_xboole_0 X0 X1) \Leftrightarrow (\forall X3.(X3 \in X2) \Leftrightarrow ((X3 \in X0) \wedge (X3 \in X1))) \tag{5}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (6)$$

Assume the following.

$$\forall X0.(v1_relat_1 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0))\Rightarrow(v1_relat_1 X1)) \quad (7)$$

Theorem 1

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((v1_funct_1 X1)\wedge(\\ m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers))))\Rightarrow(\\ \forall X2.((v1_funct_1 X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ X0 k1_numbers))))\Rightarrow((\forall X3.(X3 \in k1_relset_1 X0 X1)\Rightarrow(r1_xxreal_0 \\ (k1_seq_1 X1 X3) (k1_seq_1 X2 X3)))\Rightarrow(v6_supinf_2 (k47_valued_1 \\ X0 k1_numbers k1_numbers X2 X1)))))) \end{aligned}$$