

t422_xxreal_1 (TMVbgmAZyZdr- wozPFx6TTFPFDnVpAZ7XFhA)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k2_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xxreal_0 : \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k4_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((\neg r1_xxreal_0 X1 X0) \Rightarrow (k2_xxreal_1 X0 X1 = k2_xboole_0 (k1_tarski X0) (k4_xxreal_1 X0 X1)))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\neg (X0 \in k1_numbers) \wedge (r1_xxreal_0 X0 k2_xxreal_0)) \quad (2)$$

Assume the following.

$$v1_xxreal_0 k2_xxreal_0 \quad (3)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Leftrightarrow (X0 \in k1_numbers) \quad (4)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (v1_xxreal_0 X0) \quad (5)$$

Theorem 1

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (k2_xxreal_1 k2_xxreal_0 X0 = k2_xboole_0 (k1_tarski k2_xxreal_0) (k4_xxreal_1 k2_xxreal_0 X0))$$