

t424\_xxreal\_1

(TMcT1vHhKpajUwpruks4H4LP5yR2mPhV7me)

October 27, 2020

Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $k1\_xxreal\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xxreal\_0 : \iota$  be given. Let  $k2\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_xxreal\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_tarski : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (r1\_xxreal\_0 X0 k1\_xxreal\_0) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow ((r1\_xxreal\_0 X0 X1) \Rightarrow (k1\_xxreal\_1 X0 X1 = k2\_xboole\_0 (k4\_xxreal\_1 X0 X1) (k2\_tarski X0 X1)))) \quad (2)$$

Assume the following.

$$v1\_xxreal\_0 k1\_xxreal\_0 \quad (3)$$

**Theorem 1**

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (k1\_xxreal\_1 X0 k1\_xxreal\_0 = k2\_xboole\_0 (k4\_xxreal\_1 X0 k1\_xxreal\_0) (k2\_tarski X0 k1\_xxreal\_0))$$