

t428\_xxreal\_1  
(TMEmEdT1HbR724GskH8rXEaqxpDjohipALF)

October 27, 2020

Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $k3\_xxreal\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xxreal\_0 : \iota$  be given. Let  $k2\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_xxreal\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\neg(X0 \in k1\_numbers) \wedge (r1\_xxreal\_0 k1\_xxreal\_0 X0)) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow ((\neg r1\_xxreal\_0 X1 X0) \Rightarrow (k3\_xxreal\_1 X0 X1 = k2\_xboole\_0 (k4\_xxreal\_1 X0 X1) (k1\_tarski X1)))) \quad (2)$$

Assume the following.

$$v1\_xxreal\_0 k1\_xxreal\_0 \quad (3)$$

Assume the following.

$$\forall X0.(v1\_xxreal\_0 X0) \Leftrightarrow (X0 \in k1\_numbers) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.k2\_xboole\_0 X0 X1 = k2\_xboole\_0 X1 X0 \quad (5)$$

Assume the following.

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (v1\_xxreal\_0 X0) \quad (6)$$

**Theorem 1**

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (k3\_xxreal\_1 X0 k1\_xxreal\_0 = k2\_xboole\_0 (k4\_xxreal\_1 X0 k1\_xxreal\_0) (k1\_tarski k1\_xxreal\_0))$$