

t42_euclid_3

(TMLuHkfCAQGkrmRTCLxekna6ijXdetMCztj)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k23_rvsum_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_complex1 : \iota \Rightarrow \iota$ be given. Let $k1_complex2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_euclid_3 : \iota \Rightarrow \iota$ be given. Let $k7_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k17_euclid : \iota \Rightarrow \iota$ be given. Let $k18_euclid : \iota \Rightarrow \iota$ be given. Let $k2_numbers : \iota$ be given. Let $k4_complex1 : \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_complex1 : \iota$ be given. Let $k1_numbers : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (k23_rvsum_1 X0 X1 = k7_real_1 (k8_real_1 (k17_euclid X0) (k17_euclid \\ & X1)) (k8_real_1 (k18_euclid X0) (k18_euclid X1)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k2_numbers) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 k2_numbers) \Rightarrow (k3_complex1 (k1_complex2 X0 X1) = k7_real_1 (k8_real_1 \\ & (k3_complex1 X0) (k3_complex1 X1)) (k8_real_1 (k4_complex1 X0) \\ & (k4_complex1 X1)))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((k3_complex1 \\ & (k2_xcmplx_0 X0 (k3_xcmplx_0 X1 k7_complex1)) = X0) \wedge (k4_complex1 \\ & (k2_xcmplx_0 X0 (k3_xcmplx_0 X1 k7_complex1)) = X1))) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (m1_subset_1 (k2_euclid_3 X0) k2_numbers) \end{aligned} \tag{4}$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (m1_subset_1 (k18_euclid X0) k1_numbers) \end{aligned} \tag{5}$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow (m1_subset_1 (k17_euclid X0) k1_numbers) \quad (6)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow (k2_euclid_3 X0 = k2_xcmplx_0 (k17_euclid X0) (k3_xcmplx_0 (k18_euclid X0) k7_complex1)) \quad (7)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (8)$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2))) \Rightarrow (k23_rvsum_1 X0 X1 = k3_complex1 (k1_complex2 (k2_euclid_3 X0) (k2_euclid_3 X1))))$$