

t42_finseq_4 (TMGz-
fUsK7eXB83PdZFWgahWZc1fPYbnnUMZ)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k4_finseq_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k6_finseq_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (k4_finseq_1 X0 = k9_xtuple_0 X0) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow ((v1_relat_1 (k6_finseq_4 X0 X1)) \wedge ((v1_funct_1 (k6_finseq_4 X0 X1)) \wedge (v1_finseq_1 (k6_finseq_4 X0 X1)))) \quad (2)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (m2_subset_1 (k3_finseq_1 X0) k1_numbers k5_numbers) \quad (3)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (\forall X1. (X1 \in k10_xtuple_0 X0) \Rightarrow (\forall X2. ((v1_relat_1 X2) \wedge ((v1_funct_1 X2) \wedge (v1_finseq_1 X2)))) \Rightarrow ((X2 = k6_finseq_4 X0 X1) \Leftrightarrow ((k3_finseq_1 X2 = k6_xcmplx_0 (k3_finseq_1 X0) (k4_finseq_4 X0 X1)) \wedge (\forall X3. (v7_ordinal1 X3) \Rightarrow ((X3 \in k4_finseq_1 X2) \Rightarrow (k1_funct_1 X2 X3 = k1_funct_1 X0 (k1_nat_1 X3 (k4_finseq_4 X0 X1)))))))) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow ((X1 = k3_finseq_1 \\ & X0) \Leftrightarrow (k2_finseq_1 X1 = k9_xtuple_0 X0))) \end{aligned} \tag{5}$$

Theorem 1

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1. \forall X2.(v7_ordinal1 X2) \Rightarrow (((X1 \in k10_xtuple_0 \\ & X0) \wedge (X2 = k6_xcmplx_0 (k3_finseq_1 X0) (k4_finseq_4 X0 X1))) \Rightarrow (\\ & k4_finseq_1 (k6_finseq_4 X0 X1) = k2_finseq_1 X2))) \end{aligned}$$