

## t42\_finseq\_5

(TMTn7rC3DykKHX1pW4fh31VUVK5RJFgG7HF)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k1\_finseq\_5 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_finseq\_4 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k16\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $m1\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $k17\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge (( \\ v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \Rightarrow ((r1\_xxreal\_0 X0 (k3\_finseq\_1 \\ X1)) \Rightarrow (k3\_finseq\_1 (k16\_finseq\_1 X0 X1) = X0))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ (\forall X1.(X1 \in k10\_xtuple\_0 X0) \Rightarrow ((r1\_xxreal\_0 np\_1 (k4\_finseq\_4 \\ X0 X1)) \wedge (r1\_xxreal\_0 (k4\_finseq\_4 X0 X1) (k3\_finseq\_1 X0)))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.(m2\_finseq\_1 X1 X0) \Leftrightarrow (m1\_finseq\_1 X1 X0) \quad (3)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v7\_ordinal1 X1) \wedge (m1\_finseq\_1 \\ X2 X0)) \Rightarrow (k17\_finseq\_1 X0 X1 X2 = k16\_finseq\_1 X1 X2) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(m2\_finseq\_1 X1 X0)\Rightarrow((v1\_funct\_1 X1)\wedge((v1\_finseq\_1 X1)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers X0)))))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_1 X1 X0)\Rightarrow((v1\_relat\_1 X1)\wedge((v1\_funct\_1 X1)\wedge(v1\_finseq\_1 X1))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v1\_finseq\_1 X0)))\Rightarrow(m1\_subset\_1 (k4\_finseq\_4 X0 X1) k5\_numbers) \quad (8)$$

Assume the following.

$$\forall X0.(\neg v1\_xboole\_0 X0)\Rightarrow(\forall X1.(m2\_finseq\_1 X1 X0)\Rightarrow(\forall X2.k1\_finseq\_5 X0 X1 X2 = k17\_finseq\_1 X0 (k4\_finseq\_4 X1 X2) X1)) \quad (9)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1)\Rightarrow(v7\_ordinal1 X0) \quad (10)$$

**Theorem 1**

$$\forall X0.(\neg v1\_xboole\_0 X0)\Rightarrow(\forall X1.(m1\_subset\_1 X1 X0)\Rightarrow(\forall X2.(m2\_finseq\_1 X2 X0)\Rightarrow((X1 \in k10\_xtuple\_0 X2)\Rightarrow(k3\_finseq\_1 (k1\_finseq\_5 X0 X2 X1) = k4\_finseq\_4 X2 X1))))$$