

t42_finseq_6

(TMd46tMvtrqMEA5xWF2Hn7TNETgRCMRzSt1)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.((\neg v1_xboole_0 X0) \wedge ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0)))) \Rightarrow ((np_1 \in k4_finseq_1 X0) \wedge (k3_finseq_1 X0 \in k4_finseq_1 X0)) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (3)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (k4_finseq_1 X0 = k9_xtuple_0 X0) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0) \Rightarrow ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1) \wedge ((v5_relat_1 X1 X0) \wedge (v1_funct_1 X1))) \Rightarrow (\forall X2.(X2 \in k9_xtuple_0 X1) \Rightarrow (k7_partfun1 X0 X1 X2 = k1_funct_1 X1 X2)) \quad (6)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.(X1 = k10_xtuple_0 X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (\exists X3.(X3 \in k9_xtuple_0 X0) \wedge (X2 = k1_funct_1 X0 X3)))) \quad (7)$$

Assume the following.

$$k1_xboole_0 = the (\lambda X0 : \iota.v1_xboole_0 X0) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0) \Rightarrow (v5_relat_1 X1 X0) \quad (9)$$

Theorem 1

$$\forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(m2_finseq_1 X1 X0) \Rightarrow ((X1 \neq k1_xboole_0) \Rightarrow (k7_partfun1 X0 X1 np_1 \in k10_xtuple_0 X1)))$$