

t42_flgang_1 (TMRjrnT- nWG3TqhB82DosN4o3AuJVbrYWj1X)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k3_catalan2 : \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_flang_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_flang_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k3_catalan2 X0))) \Rightarrow ((X1 \in k8_flang_1 X0 X2) \Leftrightarrow (\exists X3. (v7_ordinal1 X3) \wedge (X1 \in k7_flang_1 X0 X2 X3))) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (2)$$

Theorem 1

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k3_catalan2 X0))) \Rightarrow (\forall X2. (v7_ordinal1 X2) \Rightarrow (r1_tarski (k7_flang_1 X0 X1 X2) (k8_flang_1 X0 X1)))$$