

t42_glib_000 (TM-
MjiFXE7XmnPWxhtns1cp7AJnFXVwPVbS8)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v1_glib_000 : \iota \Rightarrow o$ be given. Let $m1_glib_000 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k24_glib_000 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_glib_000 : \iota \Rightarrow \iota$ be given. Let $k25_glib_000 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_glib_000 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. \neg(X0 \in X1) \wedge ((m1_subset_1 X1 (k1_zfmisc_1 X2)) \wedge (v1_xboole_0 X2)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0))))) \wedge (m1_glib_000 X1 X0)) \Rightarrow (m1_subset_1 (k25_glib_000 X0 X1) (k1_zfmisc_1 (k7_glib_000 X0))) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0))))) \wedge (m1_glib_000 X1 X0)) \Rightarrow ((\neg v1_xboole_0 (k24_glib_000 X0 X1)) \wedge (m1_subset_1 (k24_glib_000 X0 X1) (k1_zfmisc_1 (k6_glib_000 X0)))) \quad (5)$$

Theorem 1

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow (\forall X1.(m1_glib_000 \\ X1 X0) \Rightarrow (\forall X2.((X2 \in k24_glib_000 X0 X1) \Rightarrow (X2 \in k6_glib_000 \\ X0)) \wedge ((X2 \in k25_glib_000 X0 X1) \Rightarrow (X2 \in k7_glib_000 X0)))) \end{aligned}$$