

## t42\_lukasi\_1

(TMM9bAsyK2qsKhQ8BqShTFgiHeTdQhsAw2U)

October 27, 2020

Let  $m1\_qc\_lang1 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_qc\_lang1 : \iota \Rightarrow \iota$  be given. Let  $k3\_cqc\_lang : \iota \Rightarrow \iota$  be given. Let  $v2\_cqc\_the1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k8\_cqc\_lang : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall X0.(m1\_qc\_lang1\ X0) \Rightarrow (\forall X1.(m2\_subset\_1\ X1\ (k9\_qc\_lang1 \\ X0)\ (k3\_cqc\_lang\ X0)) \Rightarrow (\forall X2.(m2\_subset\_1\ X2\ (k9\_qc\_lang1 \\ X0)\ (k3\_cqc\_lang\ X0)) \Rightarrow (((v2\_cqc\_the1\ X1\ X0) \wedge (v2\_cqc\_the1\ (k8\_cqc\_lang \\ X0\ X1\ X2)\ X0)) \Rightarrow (v2\_cqc\_the1\ X2\ X0)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1\_qc\_lang1\ X0) \Rightarrow (\forall X1.(m2\_subset\_1\ X1\ (k9\_qc\_lang1 \\ X0)\ (k3\_cqc\_lang\ X0)) \Rightarrow (\forall X2.(m2\_subset\_1\ X2\ (k9\_qc\_lang1 \\ X0)\ (k3\_cqc\_lang\ X0)) \Rightarrow (\forall X3.(m2\_subset\_1\ X3\ (k9\_qc\_lang1 \\ X0)\ (k3\_cqc\_lang\ X0)) \Rightarrow ((v2\_cqc\_the1\ (k8\_cqc\_lang\ X0\ X1\ X2)\ X0) \Rightarrow \\ (v2\_cqc\_the1\ (k8\_cqc\_lang\ X0\ (k8\_cqc\_lang\ X0\ X2\ X3)\ (k8\_cqc\_lang \\ X0\ X1\ X3))\ X0)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((\neg v1\_xboole\_0\ X0) \wedge ((\neg v1\_xboole\_0\ X1) \wedge \\ (m1\_subset\_1\ X1\ (k1\_zfmisc\_1\ X0)))) \Rightarrow (\forall X2.(m2\_subset\_1 \\ X2\ X0\ X1) \Leftrightarrow (m1\_subset\_1\ X2\ X1)) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.(m1\_qc\_lang1\ X0) \Rightarrow (\neg v1\_xboole\_0\ (k3\_cqc\_lang\ X0)) \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.((m1\_qc\_lang1\ X0) \wedge ((m1\_subset\_1 \\ X1\ (k3\_cqc\_lang\ X0)) \wedge (m1\_subset\_1\ X2\ (k3\_cqc\_lang\ X0)))) \Rightarrow (m2\_subset\_1 \\ (k8\_cqc\_lang\ X0\ X1\ X2)\ (k9\_qc\_lang1\ X0)\ (k3\_cqc\_lang\ X0)) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.(m1\_qc\_lang1 X0) \Rightarrow (m1\_subset\_1 (k3\_cqc\_lang X0) (k1\_zfmisc\_1 (k9\_qc\_lang1 X0))) \quad (6)$$

Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)) \Rightarrow (v1\_xboole\_0 X1)) \quad (7)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(m2\_subset\_1 X1 (k9\_qc\_lang1 \\ & X0) (k3\_cqc\_lang X0)) \Rightarrow (\forall X2.(m2\_subset\_1 X2 (k9\_qc\_lang1 \\ & X0) (k3\_cqc\_lang X0)) \Rightarrow (\forall X3.(m2\_subset\_1 X3 (k9\_qc\_lang1 \\ & X0) (k3\_cqc\_lang X0)) \Rightarrow (((v2\_cqc\_the1 (k8\_cqc\_lang X0 X1 X2) X0) \wedge \\ & (v2\_cqc\_the1 (k8\_cqc\_lang X0 X2 X3) X0)) \Rightarrow (v2\_cqc\_the1 (k8\_cqc\_lang \\ & X0 X1 X3) X0)))))) \end{aligned}$$