

t42_rewrite1
(TMKtdfxD6w2ztzhuzeGy2hSSgAEAAqqCtYoM)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $r1_rewrite1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r5_rewrite1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_relat_1 X0) \Rightarrow (\forall X1.\forall X2.\forall X3. \\ & ((r1_rewrite1 X0 X1 X2) \wedge (r1_rewrite1 X0 X2 X3)) \Rightarrow (r1_rewrite1 X0 \\ & \quad X1 X3)) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_relat_1 X0) \Rightarrow (\forall X1.\forall X2.(r5_rewrite1 \\ & X0 X1 X2) \Leftrightarrow (\exists X3.(r1_rewrite1 X0 X1 X3) \wedge (r1_rewrite1 X0 X2 \\ & \quad X3))) \end{aligned} \tag{2}$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_relat_1 X0) \Rightarrow (\forall X1.\forall X2.\forall X3. \\ & (((r1_rewrite1 X0 X1 X2) \wedge (r5_rewrite1 X0 X2 X3)) \vee ((r5_rewrite1 \\ & X0 X1 X2) \wedge (r1_rewrite1 X0 X3 X2))) \Rightarrow (r5_rewrite1 X0 X1 X3)) \end{aligned}$$