

t42_square_1 (TMQgnNUE- qJXXMVXPx2d4X1cizxsuhGdPMYc)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $np_1 : \iota$ be given. Let $k3_square_1 : \iota \Rightarrow \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 k6_numbers X0) \wedge (r1_xxreal_0 X1 np_1)) \Rightarrow (r1_xxreal_0 (k3_xcmplx_0 X0 X1) X0))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k3_square_1 X0 = k3_xcmplx_0 X0 X0) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (3)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (((r1_xxreal_0 k6_numbers X0) \wedge (r1_xxreal_0 X0 np_1)) \Rightarrow (r1_xxreal_0 (k3_square_1 X0) X0))$$