

## t42\_valuat\_1

(TMNbBYi4DREeynYvwN7UyKcs9KQEkN4SE9N)

October 27, 2020

Let  $m1\_qc\_lang1 : \iota \Rightarrow o$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_qc\_lang1 : \iota \Rightarrow \iota$  be given. Let  $k3\_cqc\_lang : \iota \Rightarrow \iota$  be given. Let  $m1\_valuat\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $r2\_valuat\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k8\_cqc\_lang : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_cqc\_lang : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $m2\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k3\_qc\_lang1 : \iota \Rightarrow \iota$  be given. Let  $k2\_valuat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_valuat\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned}
 & \forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(\neg v1\_xboole\_0 X1) \Rightarrow ( \\
 & \forall X2.(m2\_funct\_2 X2 (k3\_qc\_lang1 X0) X1 (k2\_valuat\_1 X0 X1)) \Rightarrow \\
 & (\forall X3.(m2\_subset\_1 X3 (k9\_qc\_lang1 X0) (k3\_cqc\_lang X0)) \Rightarrow \\
 & (\forall X4.(m2\_subset\_1 X4 (k9\_qc\_lang1 X0) (k3\_cqc\_lang X0)) \Rightarrow \\
 & (\forall X5.(m1\_valuat\_1 X5 X0 X1) \Rightarrow (r1\_valuat\_1 X0 X1 (k8\_cqc\_lang \\
 & X0 X3 (k8\_cqc\_lang X0 (k6\_cqc\_lang X0 X3) X4)) X5 X2))))))
 \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned}
 & \forall X0.\forall X1.((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge \\
 & (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)))) \Rightarrow (\forall X2.(m2\_subset\_1 \\
 & X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1))
 \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\neg v1\_xboole\_0 (k3\_cqc\_lang X0)) \tag{3}$$

Assume the following.

$$\begin{aligned}
 & \forall X0.\forall X1.\forall X2.((m1\_qc\_lang1 X0) \wedge ((m1\_subset\_1 \\
 & X1 (k3\_cqc\_lang X0)) \wedge (m1\_subset\_1 X2 (k3\_cqc\_lang X0)))) \Rightarrow (m2\_subset\_1 \\
 & (k8\_cqc\_lang X0 X1 X2) (k9\_qc\_lang1 X0) (k3\_cqc\_lang X0))
 \end{aligned} \tag{4}$$

Assume the following.

$$\begin{aligned}
 & \forall X0.\forall X1.((m1\_qc\_lang1 X0) \wedge (m1\_subset\_1 X1 (k3\_cqc\_lang \\
 & X0))) \Rightarrow (m2\_subset\_1 (k6\_cqc\_lang X0 X1) (k9\_qc\_lang1 X0) (k3\_cqc\_lang \\
 & X0))
 \end{aligned} \tag{5}$$

Assume the following.

$$\forall X0.(m1\_qc\_lang1 X0) \Rightarrow (m1\_subset\_1 (k3\_cqc\_lang X0) (k1\_zfmisc\_1 (k9\_qc\_lang1 X0))) \quad (6)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(\neg v1\_xboole\_0 X1) \Rightarrow ( \\ \forall X2.(m1\_valuat\_1 X2 X0 X1) \Rightarrow (\forall X3.(m2\_subset\_1 X3 \\ (k9\_qc\_lang1 X0) (k3\_cqc\_lang X0)) \Rightarrow ((r2\_valuat\_1 X0 X1 X2 X3) \Leftrightarrow \\ (\forall X4.(m2\_funct\_2 X4 (k3\_qc\_lang1 X0) X1 (k2\_valuat\_1 X0 \\ X1)) \Rightarrow (r1\_valuat\_1 X0 X1 X3 X2 X4)))))) \end{aligned} \quad (7)$$

Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)) \Rightarrow (v1\_xboole\_0 X1)) \quad (8)$$

**Theorem 1**

$$\begin{aligned} \forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(\neg v1\_xboole\_0 X1) \Rightarrow ( \\ \forall X2.(m2\_subset\_1 X2 (k9\_qc\_lang1 X0) (k3\_cqc\_lang X0)) \Rightarrow \\ (\forall X3.(m2\_subset\_1 X3 (k9\_qc\_lang1 X0) (k3\_cqc\_lang X0)) \Rightarrow \\ (\forall X4.(m1\_valuat\_1 X4 X0 X1) \Rightarrow (r2\_valuat\_1 X0 X1 X4 (k8\_cqc\_lang \\ X0 X2 (k8\_cqc\_lang X0 (k6\_cqc\_lang X0 X2) X3)))))) \end{aligned}$$