

t43_bvfunc14 (TM-
MusYSLrcR6hwwczVQ6Q7B1WA7SGUz5HN5)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k1_bvfunc_2 : \iota \Rightarrow \iota$ be given. Let $m1_eqrel_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_enumset1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_bvfunc_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_partit1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_tarSKI : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_enumset1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 \\
 & \quad (k1_bvfunc_2 X0))) \Rightarrow (\forall X2. (m1_eqrel_1 X2 X0) \Rightarrow (\forall X3. \\
 & \quad (m1_eqrel_1 X3 X0) \Rightarrow (\forall X4. (m1_eqrel_1 X4 X0) \Rightarrow (\forall X5. \\
 & \quad (m1_eqrel_1 X5 X0) \Rightarrow (\forall X6. (m1_eqrel_1 X6 X0) \Rightarrow (\forall X7. \\
 & \quad (m1_eqrel_1 X7 X0) \Rightarrow (\forall X8. (m1_eqrel_1 X8 X0) \Rightarrow ((X1 = k5_enumset1 \\
 & \quad X2 X3 X4 X5 X6 X7 X8) \Rightarrow ((X2 = X3) \vee ((X2 = X4) \vee ((X2 = X5) \vee ((X2 = X6) \vee ((X2 = \\
 & \quad X7) \vee ((X2 = X8) \vee ((X3 = X4) \vee ((X3 = X5) \vee ((X3 = X6) \vee ((X3 = X7) \vee ((X3 = X8) \vee \\
 & \quad ((X4 = X5) \vee ((X4 = X6) \vee ((X4 = X7) \vee ((X4 = X8) \vee ((X5 = X6) \vee ((X5 = X7) \vee (\\
 & \quad (X5 = X8) \vee ((X6 = X7) \vee ((X6 = X8) \vee ((X7 = X8) \vee (k5_bvfunc_2 X0 X2 X1 = k2_partit1 \\
 & \quad X0 (k2_partit1 X0 (k2_partit1 X0 (k2_partit1 X0 (k2_partit1 X0 X3 \\
 & \quad X4) X5) X6) X7) X8)))))))))))))))))))))))))))))))))) \\
 & \hspace{15em} (1)
 \end{aligned}$$

Assume the following.

$$\begin{aligned}
 & \forall X0. \forall X1. \forall X2. \forall X3. \forall X4. \forall X5. \\
 & \forall X6. k5_enumset1 X0 X1 X2 X3 X4 X5 X6 = k2_xboole_0 (k2_tarSKI \\
 & \quad X0 X1) (k3_enumset1 X2 X3 X4 X5 X6) \\
 & \hspace{15em} (2)
 \end{aligned}$$

Assume the following.

$$\forall X0. \forall X1. k2_tarSKI X0 X1 = k2_tarSKI X1 X0 \hspace{15em} (3)$$

Theorem 1

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 \\ (k1_bfunc_2 X0))) \Rightarrow (\forall X2.(m1_eqrel_1 X2 X0) \Rightarrow (\forall X3. \\ (m1_eqrel_1 X3 X0) \Rightarrow (\forall X4.(m1_eqrel_1 X4 X0) \Rightarrow (\forall X5. \\ (m1_eqrel_1 X5 X0) \Rightarrow (\forall X6.(m1_eqrel_1 X6 X0) \Rightarrow (\forall X7. \\ (m1_eqrel_1 X7 X0) \Rightarrow (\forall X8.(m1_eqrel_1 X8 X0) \Rightarrow ((X1 = k5_enumset1 \\ X2 X3 X4 X5 X6 X7 X8) \Rightarrow ((X2 = X3) \vee ((X2 = X4) \vee ((X2 = X5) \vee ((X2 = X6) \vee ((X2 = \\ X7) \vee ((X2 = X8) \vee ((X3 = X4) \vee ((X3 = X5) \vee ((X3 = X6) \vee ((X3 = X7) \vee ((X3 = X8) \vee \\ ((X4 = X5) \vee ((X4 = X6) \vee ((X4 = X7) \vee ((X4 = X8) \vee ((X5 = X6) \vee ((X5 = X7) \vee (\\ (X5 = X8) \vee ((X6 = X7) \vee ((X6 = X8) \vee ((X7 = X8) \vee (k5_bfunc_2 X0 X3 X1 = k2_partit1 \\ X0 (k2_partit1 X0 (k2_partit1 X0 (k2_partit1 X0 (k2_partit1 X0 X2 \\ X4) X5) X6) X7) X8))))))))))))))))))))))))))))))))))))))))))))))))))))))))))))) \end{aligned}$$