

t43_fcont_1
(TMJPX7ntMCvdTyifzG2gJAKabZ1mHZxik8g)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_square_1 : \iota \Rightarrow \iota$ be given. Let $v1_fcont_1 : \iota \Rightarrow o$ be given. Let $k2_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow (k9_xtuple_0 (k5_relat_1 X1 X0) = k3_xboole_0 (k9_xtuple_0 X1) X0) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((X0 \in k9_xtuple_0 (k5_relat_1 X2 X1)) \Rightarrow (k1_funct_1 (k5_relat_1 X2 X1) X0 = k1_funct_1 X2 X0)) \quad (2)$$

Assume the following.

$$\forall X0. ((v1_funct_1 X0) \wedge (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k1_numbers k1_numbers)))) \Rightarrow ((\forall X1. (v1_xreal_0 X1) \Rightarrow ((X1 \in k1_relset_1 k1_numbers X0) \Rightarrow (k1_seq_1 X0 X1 = k3_square_1 X1))) \Rightarrow (v1_fcont_1 (k2_partfun1 k1_numbers k1_numbers X0 (k1_relset_1 k1_numbers X0)))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Rightarrow (k3_xboole_0 X0 X1 = X0) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((v1_funct_1 X2)\wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow(k2_partfun1 X0 X1 X2 X3 = k5_relat_1 X2 X3) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v3_valued_0 X0)))\Rightarrow(k1_seq_1 X0 X1 = k1_funct_1 X0 X1) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (7)$$

Assume the following.

$$\forall X0.(v1_relat_1 X0)\Rightarrow(k5_relat_1 X0 (k9_xtuple_0 X0) = X0) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1_relat_1 X2)\wedge(v4_relat_1 X2 X1))\Rightarrow((v1_relat_1 (k5_relat_1 X2 X0))\wedge((v4_relat_1 (k5_relat_1 X2 X0) X0)\wedge(v4_relat_1 (k5_relat_1 X2 X0) X1))) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((v1_funct_1 X2)\wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow((v1_funct_1 (k2_partfun1 X0 X1 X2 X3))\wedge(m1_subset_1 (k2_partfun1 X0 X1 X2 X3) (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.k3_xboole_0 X0 X1 = k3_xboole_0 X1 X0 \quad (11)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v5_relat_1 X0 k1_numbers))\Rightarrow((v1_relat_1 X0)\wedge(v3_valued_0 X0)) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (14)$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. ((v1_funct_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (k2_zfmisc_1 k1_numbers k1_numbers)))) \Rightarrow (((r1_tarski X0 (k1_relset_1 \\ & k1_numbers X1)) \wedge (\forall X2. (v1_xreal_0 X2) \Rightarrow ((X2 \in X0) \Rightarrow (k1_seq_1 \\ & X1 X2 = k3_square_1 X2)))) \Rightarrow (v1_fcont_1 (k2_partfun1 k1_numbers \\ & k1_numbers X1 X0))) \end{aligned}$$