

t43_finseq_5 (TMKr-
jpxJi4HBDwAyvx5A4XHTQrEfAVMGH7H)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k4_finseq_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_finseq_5 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k17_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m2_finseq_1 X1 X0) \Rightarrow (\forall X2. (v7_ordinal1 \\ & X2) \Rightarrow (\forall X3. (v7_ordinal1 X3) \Rightarrow (((X2 \in k4_finseq_1 X1) \wedge (X3 \in \\ & k2_finseq_1 X2)) \Rightarrow ((X3 \in k4_finseq_1 X1) \wedge (k7_partfun1 X0 (k17_finseq_1 \\ & X0 X2 X1) X3 = k7_partfun1 X0 X1 X3)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1. (X1 \in k10_xtuple_0 X0) \Rightarrow (k4_finseq_4 X0 X1 \in k4_finseq_1 \\ & X0)) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (3)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m2_finseq_1 X1 X0) \Rightarrow ((v1_funct_1 X1) \wedge (\\ & (v1_finseq_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers \\ & X0)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow((v1_relat_1 X1)\wedge(v1_funct_1 X1)\wedge(v1_finseq_1 X1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))\Rightarrow(m1_subset_1 (k4_finseq_4 X0 X1) k5_numbers) \quad (7)$$

Assume the following.

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.(m2_finseq_1 X1 X0)\Rightarrow(\forall X2.k1_finseq_5 X0 X1 X2 = k17_finseq_1 X0 (k4_finseq_4 X1 X2) X1)) \quad (8)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (9)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(\forall X1.(\neg v1_xboole_0 X1)\Rightarrow(\forall X2.(m1_subset_1 X2 X1)\Rightarrow(\forall X3.(m2_finseq_1 X3 X1)\Rightarrow(((X2 \in k10_xtuple_0 X3)\wedge(X0 \in k2_finseq_1 (k4_finseq_4 X3 X2))))\Rightarrow(k7_partfun1 X1 (k1_finseq_5 X1 X3 X2) X0 = k7_partfun1 X1 X3 X0))))))$$