

t43_fomodel0

(TMWV7ZAeHYmgGeMsyKbZuBtFG8p11sCnxoN)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k16_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.(r1_partfun1 (k7_funcop_1 X0 X2) (k7_funcop_1 X1 X3)) \Leftrightarrow ((X2 = X3) \vee (r1_xboole_0 X0 X1)) \quad (1)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_partfun1 X0 X1) \Leftrightarrow (k1_funct_4 X0 X1 = k1_funct_4 X1 X0))) \quad (2)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_tarski (k9_xtuple_0 X0) (k9_xtuple_0 X1)) \Rightarrow (k1_funct_4 X0 X1 = X1))) \quad (3)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2.((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow (k1_funct_4 (k1_funct_4 X0 X1) X2 = k1_funct_4 X0 (k1_funct_4 X1 X2)))) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.(k9_xtuple_0 (k2_funcop_1 X0 X1) = X0) \wedge (r1_tarski (k10_xtuple_0 (k2_funcop_1 X0 X1)) (k1_tarski X1)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(X0\neq X1)\Rightarrow(r1_xboole_0 (k1_tarski X0) (k1_tarski X1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.r1_tarski X0 X0 \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.k7_funcop_1 X0 X1 = k2_funcop_1 X0 X1 \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.v1_relat_1 (k2_zfmisc_1 X0 X1) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.(v1_funct_1 (k7_funcop_1 X0 X1))\wedge((v1_funct_2 (k7_funcop_1 X0 X1) X0 (k1_tarski X1))\wedge(m1_subset_1 (k7_funcop_1 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 (k1_tarski X1)))))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.k16_funcop_1 X0 X1 = k7_funcop_1 (k1_tarski X0) X1 \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.k2_funcop_1 X0 X1 = k2_zfmisc_1 X0 (k1_tarski X1) \quad (12)$$

Theorem 1

$$\forall X0.\forall X1.\forall X2.\forall X3.\forall X4.((v1_relat_1 X4)\wedge(v1_funct_1 X4))\Rightarrow(((X0 = X1)\Rightarrow(k1_funct_4 (k1_funct_4 X4 (k16_funcop_1 X0 X2)) (k16_funcop_1 X1 X3) = k1_funct_4 X4 (k16_funcop_1 X1 X3)))\wedge((X0\neq X1)\Rightarrow(k1_funct_4 (k1_funct_4 X4 (k16_funcop_1 X0 X2)) (k16_funcop_1 X1 X3) = k1_funct_4 (k1_funct_4 X4 (k16_funcop_1 X1 X3)) (k16_funcop_1 X0 X2))))$$