

# t43\_lpspace2

(TMdrbChWJetzTX4Ax142hnWHkhJjrKZbZjK)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v1\_prob\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v4\_prob\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k7\_numbers : \iota$  be given. Let  $v10\_valued\_0 : \iota \Rightarrow o$  be given. Let  $v6\_supinf\_2 : \iota \Rightarrow o$  be given. Let  $v4\_measure1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_numbers : \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $k1\_lpspace2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_lpspace2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_lpspace1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned}
& \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.((\neg v1\_xboole\_0 X1) \wedge \\
& ((v1\_prob\_1 X1 X0) \wedge ((v4\_prob\_1 X1 X0) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 \\
& (k1\_zfmisc\_1 X0)))))) \Rightarrow (\forall X2.((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 \\
& X2 X1 k7\_numbers) \wedge ((v10\_valued\_0 X2) \wedge ((v6\_supinf\_2 X2) \wedge ((v4\_measure1 \\
& X2 X0 X1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X1 k7\_numbers))))))) \Rightarrow \\
& (\forall X3.((v1\_funct\_1 X3) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\
& X0 k1\_numbers)))) \Rightarrow (\forall X4.((v1\_funct\_1 X4) \wedge (m1\_subset\_1 \\
& X4 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k1\_numbers)))) \Rightarrow (\forall X5.( \\
& (v2\_xxreal\_0 X5) \wedge (m1\_subset\_1 X5 k1\_numbers)) \Rightarrow ((r1\_lpspace1 \\
& X0 X1 X2 X3 X4) \Rightarrow (k5\_lpspace2 X0 X1 X2 X3 X5 = k5\_lpspace2 X0 X1 X2 X4 X5))))))
\end{aligned}$$

(1)

Assume the following.

$$\begin{aligned}
& \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.((\neg v1\_xboole\_0 X1) \wedge \\
& ((v1\_prob\_1 X1 X0) \wedge ((v4\_prob\_1 X1 X0) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 \\
& (k1\_zfmisc\_1 X0)))))) \Rightarrow (\forall X2.((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 \\
& X2 X1 k7\_numbers) \wedge ((v10\_valued\_0 X2) \wedge ((v6\_supinf\_2 X2) \wedge ((v4\_measure1 \\
& X2 X0 X1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X1 k7\_numbers)))))) \Rightarrow \\
& (\forall X3.((v1\_funct\_1 X3) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\
& X0 k1\_numbers)))) \Rightarrow (\forall X4.((v2\_xxreal\_0 X4) \wedge (m1\_subset\_1 \\
& X4 k1\_numbers)) \Rightarrow (k5\_lpspace2 X0 X1 X2 X3 X4 = ReplSep (toset (\lambda X5 : \\
& \iota.(v1\_funct\_1 X5) \wedge (m1\_subset\_1 X5 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\
& X0 k1\_numbers)))))) (\lambda X5 : \iota.(X5 \in k1\_lpspace2 X0 X1 X2 X4) \wedge ( \\
& r1\_lpspace1 X0 X1 X2 X3 X5)) (\lambda X5 : \iota.X5))))))
\end{aligned} \tag{2}$$

**Theorem 1**

$$\begin{aligned}
& \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.((\neg v1\_xboole\_0 X1) \wedge \\
& ((v1\_prob\_1 X1 X0) \wedge ((v4\_prob\_1 X1 X0) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 \\
& (k1\_zfmisc\_1 X0)))))) \Rightarrow (\forall X2.((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 \\
& X2 X1 k7\_numbers) \wedge ((v10\_valued\_0 X2) \wedge ((v6\_supinf\_2 X2) \wedge ((v4\_measure1 \\
& X2 X0 X1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X1 k7\_numbers)))))) \Rightarrow \\
& (\forall X3.((v1\_funct\_1 X3) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\
& X0 k1\_numbers)))) \Rightarrow (\forall X4.((v1\_funct\_1 X4) \wedge (m1\_subset\_1 \\
& X4 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k1\_numbers)))) \Rightarrow (\forall X5.( \\
& (v2\_xxreal\_0 X5) \wedge (m1\_subset\_1 X5 k1\_numbers)) \Rightarrow (((X3 \in k1\_lpspace2 \\
& X0 X1 X2 X5) \wedge (X4 \in k5\_lpspace2 X0 X1 X2 X3 X5)) \Rightarrow (k5\_lpspace2 X0 X1 X2 \\
& X3 X5 = k5\_lpspace2 X0 X1 X2 X4 X5))))))
\end{aligned}$$