

t43_partfun1

(TMHJuzmQXd6URR41JiuGLZbmefYN5YK6cDd)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((v1_relat_1 X3) \wedge \\ & (v1_funct_1 X3)) \Rightarrow ((X0 \in k1_relset_1 X1 (k3_partfun1 X3 X1 X2)) \Leftrightarrow \\ & ((X0 \in k9_xtuple_0 X3) \wedge ((X0 \in X1) \wedge (k1_funct_1 X3 X0 \in X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.(v1_relat_1 X1) \Rightarrow (v1_relat_1 (k6_relat_1 X0 X1)) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.(v1_relat_1 X0) \Rightarrow (v1_relat_1 (k5_relat_1 X0 X1)) \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((v1_relat_1 X0) \wedge (v1_funct_1 \\ & X0)) \Rightarrow ((v1_funct_1 (k3_partfun1 X0 X1 X2)) \wedge (m1_subset_1 (k3_partfun1 \\ & X0 X1 X2) (k1_zfmisc_1 (k2_zfmisc_1 X1 X2)))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.\forall X2. \\ & (X2 = k7_relat_1 X0 X1) \Leftrightarrow (\forall X3.(X3 \in X2) \Leftrightarrow (\exists X4.(X4 \in k9_xtuple_0 \\ & X0) \wedge ((X4 \in X1) \wedge (X3 = k1_funct_1 X0 X4)))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(r1_tarSKI X0 X1)\Leftrightarrow(\forall X2.(X2 \in X0)\Rightarrow (X2 \in X1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow (v1_partfun1 X1 X0)\Leftrightarrow(k1_relset_1 X0 X1 = X0) \quad (7)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v1_funct_1 X0))\Rightarrow(\forall X1.\forall X2. k3_partfun1 X0 X1 X2 = k5_relat_1 (k6_relat_1 X2 X0) X1) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (9)$$

Theorem 1

$$\forall X0.\forall X1.\forall X2.((v1_relat_1 X2)\wedge(v1_funct_1 X2))\Rightarrow((v1_partfun1 (k3_partfun1 X2 X0 X1) X0)\Rightarrow(r1_tarSKI (k7_relat_1 X2 X0) X1))$$