

t43_tops_1
(TMGXTSwFsFr5wgfLxRSiqPUq2ZWdo7kVe6V)

October 27, 2020

Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $v4_pre_topc : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_tops_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tops_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k2_pre_topc : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. k4_xboole_0 X0 (k3_xboole_0 X1 X2) = k2_xboole_0 (k4_xboole_0 X0 X1) (k4_xboole_0 X0 X2) \quad (1)$$

Assume the following.

$$\forall X0. k4_xboole_0 X0 k1_xboole_0 = X0 \quad (2)$$

Assume the following.

$$\forall X0. (l1_pre_topc X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (k2_pre_topc X0 X1 = k4_subset_1 (u1_struct_0 X0) X1 (k2_tops_1 X0 X1))) \quad (3)$$

Assume the following.

$$\forall X0. k3_xboole_0 X0 k1_xboole_0 = k1_xboole_0 \quad (4)$$

Assume the following.

$$\forall X0. (l1_pre_topc X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (((v4_pre_topc X1 X0) \Rightarrow (k2_pre_topc X0 X1 = X1)) \wedge (((v2_pre_topc X0) \wedge (k2_pre_topc X0 X1 = X1)) \Rightarrow (v4_pre_topc X1 X0)))) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (k7_subset_1 X0 X1 X2 = k4_xboole_0 X1 X2) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1_subset_1 X1 (k1_zfmisc_1 X0))\wedge(m1_subset_1 X2 (k1_zfmisc_1 X0)))\Rightarrow(k4_subset_1 X0 X1 X2 = k2_xboole_0 X1 X2) \quad (7)$$

Assume the following.

$$\forall X0.(l1_pre_topc X0)\Rightarrow(\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0)))\Rightarrow(k2_tops_1 X0 X1 = k7_subset_1 (u1_struct_0 X0) (k2_pre_topc X0 X1) (k1_tops_1 X0 X1))) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((l1_pre_topc X0)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))))\Rightarrow(m1_subset_1 (k2_tops_1 X0 X1) (k1_zfmisc_1 (u1_struct_0 X0))) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.k2_xboole_0 X0 X1 = k2_xboole_0 X1 X0 \quad (10)$$

Theorem 1

$$\forall X0.((v2_pre_topc X0)\wedge(l1_pre_topc X0))\Rightarrow(\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0)))\Rightarrow((v4_pre_topc X1 X0)\Leftrightarrow(k2_tops_1 X0 X1 = k7_subset_1 (u1_struct_0 X0) X1 (k1_tops_1 X0 X1))))$$