

t43_zf_lang1
(TMMB9nEy1FkvKyVToSwyiApdtrsfwFzP8L9)

October 27, 2020

Let $v1_zf_lang : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $r3_zf_lang : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r2_zf_lang : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. ((v1_zf_lang X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow (\forall X1. \\ & ((v1_zf_lang X1) \wedge (m2_finseq_1 X1 k5_numbers)) \Rightarrow (\neg (r3_zf_lang \\ & X0 X1) \wedge (r2_zf_lang X1 X0))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_zf_lang X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow (\forall X1. \\ & ((v1_zf_lang X1) \wedge (m2_finseq_1 X1 k5_numbers)) \Rightarrow ((r3_zf_lang \\ & X0 X1) \Leftrightarrow ((r2_zf_lang X0 X1) \wedge (X0 \neq X1)))) \end{aligned} \quad (2)$$

Theorem 1

$$\begin{aligned} & \forall X0. ((v1_zf_lang X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow (\forall X1. \\ & ((v1_zf_lang X1) \wedge (m2_finseq_1 X1 k5_numbers)) \Rightarrow (\neg (r3_zf_lang \\ & X0 X1) \wedge (r3_zf_lang X1 X0))) \end{aligned}$$