

t44_complex1 (TMT- VAn7WDpqahrHuPiYYvCypAuRUjQpjql1)

October 27, 2020

Let $k17_complex1 : \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (r1_xxreal_0 k6_numbers (k3_xcmplx_0 X0 X0)) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((r1_xxreal_0 k6_numbers X0) \Rightarrow (k17_complex1 X0 = X0)) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xxreal_0 X0 k6_numbers) \wedge ((\neg r1_xxreal_0 k6_numbers X1) \wedge (r1_xxreal_0 k6_numbers (k3_xcmplx_0 X0 X1)))))) \quad (3)$$

Assume the following.

$$m1_subset_1 k6_numbers k1_numbers \quad (4)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (5)$$

Theorem 1 $k17_complex1 k6_numbers = k6_numbers$.