

t44_finseq_5 (TMcpjqSYyE- grtG8a6WEDHUnK3k5YUrvPsLZ)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_finseq_5 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k4_finseq_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Leftrightarrow (r1_tarski (k2_finseq_1 X0) (k2_finseq_1 X1)))) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(\neg v1_xboole_0 X1) \Rightarrow (\\ & \forall X2.(m1_subset_1 X2 X1) \Rightarrow (\forall X3.(m2_finseq_1 X3 X1) \Rightarrow \\ & (((X2 \in k10_xtuple_0 X3) \wedge (X0 \in k2_finseq_1 (k4_finseq_4 X3 X2))) \Rightarrow \\ & (k7_partfun1 X1 (k1_finseq_5 X1 X3 X2) X0 = k7_partfun1 X1 X3 X0)))))) \quad (2) \end{aligned}$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1.(X1 \in k10_xtuple_0 X0) \Rightarrow ((r1_xxreal_0 np_1 (k4_finseq_4 X0 X1)) \wedge (r1_xxreal_0 (k4_finseq_4 X0 X1) (k3_finseq_1 X0)))) \quad (3) \end{aligned}$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((X0 \in k2_finseq_1 X1) \Leftrightarrow ((r1_xxreal_0 np_1 X0) \wedge (r1_xxreal_0 X0 X1)))) \quad (4)$$

Assume the following.

$$\begin{aligned} & ((v2_xreal_0 \ np_1) \wedge (m2_subset_1 \ np_1 \ k1_numbers \ k5_numbers)) \wedge \\ & ((m1_subset_1 \ np_1 \ k5_numbers) \wedge (m1_subset_1 \ np_1 \ k1_numbers)) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. r1_tarski \ X0 \ X0 \quad (6)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 \ X1 \ X0) \Leftrightarrow (m1_finseq_1 \ X1 \ X0) \quad (7)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m2_finseq_1 \ X1 \ X0) \Rightarrow ((v1_funct_1 \ X1) \wedge \\ & (v1_finseq_1 \ X1) \wedge (m1_subset_1 \ X1 \ (k1_zfmisc_1 \ (k2_zfmisc_1 \ k5_numbers \\ & \ X0)))) \end{aligned} \quad (9)$$

Assume the following.

$$\forall X0. \forall X1. (m1_finseq_1 \ X1 \ X0) \Rightarrow ((v1_relat_1 \ X1) \wedge (v1_funct_1 \ X1) \wedge (v1_finseq_1 \ X1)) \quad (10)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 \ X0) \wedge ((v1_funct_1 \ X0) \wedge (v1_finseq_1 \ X0))) \Rightarrow (m1_subset_1 \ (k4_finseq_4 \ X0 \ X1) \ k5_numbers) \quad (11)$$

Assume the following.

$$\forall X0. (m1_subset_1 \ X0 \ k4_ordinal1) \Rightarrow (v7_ordinal1 \ X0) \quad (12)$$

Theorem 1

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 \ X0) \Rightarrow (\forall X1. (m1_subset_1 \ X1 \ X0) \Rightarrow \\ & (\forall X2. (m2_finseq_1 \ X2 \ X0) \Rightarrow ((X1 \in k10_xtuple_0 \ X2) \Rightarrow (k7_partfun1 \\ & \ X0 \ (k1_finseq_5 \ X0 \ X2 \ X1) \ np_1 = k7_partfun1 \ X0 \ X2 \ np_1)))) \end{aligned}$$