

t44_mssubfam (TMPBFQLoHmdhzLaHT-
piyVuRZLcqw2RP3p5d)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m3_pboole : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_mboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_pboole : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_pboole : \iota \Rightarrow \iota$ be given. Let $r6_pboole : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_mssubfam : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_pboole : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v4_relat_1 X1 X0) \wedge \\ & (v1_funct_1 X1) \wedge (v1_partfun1 X1 X0))) \Rightarrow ((r2_pboole X0 X1 (k1_pboole \\ & X0)) \Rightarrow (r6_pboole X0 X1 (k1_pboole X0))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v4_relat_1 X1 X0) \wedge \\ & (v1_funct_1 X1) \wedge (v1_partfun1 X1 X0))) \Rightarrow (\forall X2. ((v1_relat_1 \\ & X2) \wedge ((v4_relat_1 X2 X0) \wedge ((v1_funct_1 X2) \wedge (v1_partfun1 X2 X0)))) \Rightarrow \\ & (\forall X3. (m3_pboole X3 X0 (k1_mboolean X0 X1)) \Rightarrow ((r1_pboole \\ & X0 X2 X3) \Rightarrow (r2_pboole X0 (k4_mssubfam X0 X1 X3) X2)))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v4_relat_1 X1 X0) \wedge \\ & (v1_funct_1 X1) \wedge (v1_partfun1 X1 X0))) \Rightarrow (\forall X2. (m3_pboole \\ & X2 X0 X1) \Rightarrow ((v1_relat_1 X2) \wedge ((v4_relat_1 X2 X0) \wedge ((v1_funct_1 X2) \wedge \\ & (v1_partfun1 X2 X0)))))) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (((v1_relat_1 X1) \wedge ((v4_relat_1 \\ & X1 X0) \wedge ((v1_funct_1 X1) \wedge (v1_partfun1 X1 X0)))) \wedge (m3_pboole X2 \\ & X0 (k1_mboolean X0 X1))) \Rightarrow (m3_pboole (k4_mssubfam X0 X1 X2) X0 X1) \end{aligned} \tag{4}$$

Assume the following.

$$\forall X0.(v1_relat_1 (k1_pboole X0)) \wedge ((v4_relat_1 (k1_pboole X0) X0) \wedge ((v1_funct_1 (k1_pboole X0)) \wedge (v1_partfun1 (k1_pboole X0) X0))) \quad (5)$$

Theorem 1

$$\forall X0.\forall X1.((v1_relat_1 X1) \wedge ((v4_relat_1 X1 X0) \wedge (v1_funct_1 X1) \wedge (v1_partfun1 X1 X0))) \Rightarrow (\forall X2.(m3_pboole X2 X0 (k1_mboolean X0 X1)) \Rightarrow ((r1_pboole X0 (k1_pboole X0) X2) \Rightarrow (r6_pboole X0 (k4_mssubfam X0 X1 X2) (k1_pboole X0))))$$