

t44_rfunct_2 (TMX- ACVV3HdDTXVc2oSZEcHGSJZXAJnszvGF)

October 27, 2020

Let $v1_funct.1 : \iota \Rightarrow o$ be given. Let $m1_subset.1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc.1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc.1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $v6_valued.0 : \iota \Rightarrow o$ be given. Let $k2_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k9_subset.1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_relset.1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xxreal.0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_seq.1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xxreal.0 : \iota \Rightarrow o$ be given. Let $k3_xboole.0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat.1 : \iota \Rightarrow o$ be given. Let $v4_relat.1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xreal.0 : \iota \Rightarrow o$ be given. Let $v5_relat.1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_funct.1 X1) \wedge (m1_subset.1 X1 (k1_zfmisc.1 \\ & (k2_zfmisc.1 k1_numbers k1_numbers)))) \Rightarrow ((v6_valued.0 (k2_partfun1 \\ & k1_numbers k1_numbers X1 X0)) \Leftrightarrow (\forall X2. (m1_subset.1 X2 k1_numbers) \Rightarrow \\ & (\forall X3. (m1_subset.1 X3 k1_numbers) \Rightarrow (\neg (X2 \in k9_subset.1 k1_numbers \\ & X0 (k1_relset.1 k1_numbers X1)) \wedge ((X3 \in k9_subset.1 k1_numbers \\ & X0 (k1_relset.1 k1_numbers X1)) \wedge (\neg r1_xxreal.0 X3 X2) \wedge (r1_xxreal.0 \\ & (k1_seq.1 X1 X2) (k1_seq.1 X1 X3))))))))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. ((v1_xxreal.0 X0) \wedge (v1_xxreal.0 X1)) \Rightarrow (r1_xxreal.0 X0 X0) \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset.1 X2 (k1_zfmisc.1 X0)) \Rightarrow (k9_subset.1 X0 X1 X2 = k3_xboole.0 X1 X2) \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat.1 X1) \wedge (v4_relat.1 X1 X0)) \Rightarrow (m1_subset.1 (k1_relset.1 X0 X1) (k1_zfmisc.1 X0)) \tag{4}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k3_xboole.0 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in X0) \wedge (X3 \in X1))) \tag{5}$$

Assume the following.

$$\forall X0.\forall X1.(X1 = k1_tarSKI X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (6)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xxreal_0 X0) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \quad (8)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \quad (10)$$

Theorem 1

$$\forall X0.\forall X1.((v1_funct_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k1_numbers k1_numbers)))) \Rightarrow (v6_valued_0 (k2_partfun1 k1_numbers k1_numbers X1 (k1_tarSKI X0)))$$